

Immunisation Passport

for babies born on or after
1st October 2024



Name

DOB

This is an important document, please keep it safe.



Introduction

The aim of this Immunisation Passport is to ensure you have a record of all your immunisations.

These records should include all

- primary childhood immunisations,
- school immunisations,
- seasonal vaccines
- other vaccines

There are times when you may require other vaccines following an outbreak or for work or travel reasons. You can also record those vaccines in this booklet.

Please bring this Immunisation Passport with you every time you attend an immunisation appointment.

Name

Also known as

Date of Birth

Address

Contact Phone
Number(s)

Email Address

Allergies

GP Name

GP Address

GP Phone
Number

GP Name

GP Address

GP Phone
Number

GP Name

GP Address

GP Phone
Number

Please keep this booklet safe and bring it with you to every vaccination appointment.

The table below shows at what age the immunisations are given, where they are given and which vaccines are given.

When	Where	Vaccination
2 months	GP Surgery Visit 1	6 in 1+PCV+MenB+Rotavirus 3 Injections+Oral Drops
4 months	GP Surgery Visit 2	6 in 1+MenB+Rotavirus 2 Injections+Oral Drops
6 months	GP Surgery Visit 3	6 in 1+PCV 2 Injections
— No Rotavirus vaccine on or after 8 months 0 days —		
12 months	GP Surgery Visit 4	MMR+MenB+Chickenpox 3 Injections
13 months	GP Surgery Visit 5	6 in 1+MenC+PCV 3 Injections



Junior infants (first year of primary school)	School*	4 in 1 + MMRV
First year of second level school	School	HPV+Tdap+ MenACWY
Every year (Seasonal Vaccines)	GP Surgery or Pharmacy	Flu**

* These vaccines are only available in GP surgeries in Sligo, Leitrim and Donegal.

** For children aged 2 years and older and other eligible age groups and at-risk groups

6 in 1	Diphtheria, Haemophilus influenzae b (Hib), Hepatitis B, Pertussis (whooping cough), Polio, Tetanus
PCV	Pneumococcal conjugate
MenB	Meningococcal B
Rota	Rotavirus
MenC	Meningococcal C
MMR	Measles, Mumps, Rubella
Chickenpox	Varicella
MMRV	Measles, Mumps, Rubella, Chickenpox (Varicella)
4 in 1	Diphtheria, Pertussis (whooping cough), Polio, Tetanus
HPV	Human papillomavirus
Tdap	Tetanus, low dose diphtheria and acellular pertussis (whooping cough)
MenACWY	Meningococcal types A, C, W and Y
Flu	Influenza

Primary Childhood Immunisations

Vaccine Name Age Given Manufacturer Batch Number

GP Visit 1 | 2 months

6 in 1			
PCV			
MenB			
Rota			

GP Visit 2 | 4 months

Next appointment date:

6 in 1			
MenB			
Rota			

GP Visit 3 | 6 months

Next appointment date:

6 in 1			
PCV			

6 in 1 Diphtheria, Haemophilus influenzae b (Hib), Hepatitis B, Pertussis (whooping cough), Polio, Tetanus

Name

DOB

Expiry Date

Route/Site

Administered by

Date



Expiry Date	Route/Site	Administered by	Date



Expiry Date	Route/Site	Administered by	Date



Expiry Date	Route/Site	Administered by	Date

PCV Pneumococcal conjugate

MenB Meningococcal B

Rota Rotavirus

Primary Childhood Immunisations

Vaccine Name Age Given Manufacturer Batch Number

GP Visit 4 | 12 months

Next appointment date:

MMR			
MenB			
Chickenpox			

GP Visit 5 | 13 months

Next appointment date:

6 in 1			
MenC			
PCV			

Other

MMR Measles, Mumps, Rubella

MenB Meningococcal B

PCV Pneumococcal conjugate

Name


DOB

Expiry Date


Route/Site

Administered by


Date



Expiry Date	Route/Site	Administered by	Date



Expiry Date	Route/Site	Administered by	Date



Expiry Date	Route/Site	Administered by	Date

Chickenpox Varicella

6 in 1

Diphtheria, Haemophilus influenzae b (Hib), Hepatitis B, Pertussis (whooping cough), Polio, Tetanus

School Immunisations

Vaccine Name Age Given Manufacturer Batch Number

Primary School

4 in 1			
MMRV			

Secondary School

Tdap			
MenACWY			
HPV			

Other

4 in 1 Diphtheria, Pertussis (whooping cough), Polio, Tetanus

MMRV Measles Mumps Rubella and Chickenpox (Varicella)

MMR Measles, Mumps, Rubella

Tdap Tetanus, low dose diphtheria and acellular pertussis (whooping cough)

Name

DOB

Expiry Date

Route/Site

Administered by

Date



Expiry Date	Route/Site	Administered by	Date



Expiry Date	Route/Site	Administered by	Date



Expiry Date	Route/Site	Administered by	Date

MenACWY Meningococcal types A, C, W and Y

HPV Human papillomavirus

Post Vaccination Advice

What common reactions can my child get after being vaccinated and what should I do?

Common reaction	What to do
<p>A fever is common after MenB vaccine at 2 and 4 months (Visits 1 and 2).</p> <p>You should only give paracetamol to your baby after the 2 and 4 month visits</p>	<p>Give liquid infant paracetamol</p> <ol style="list-style-type: none">1. Give 2.5 mls (60 mg) at the time of the immunisation or shortly after.2. Give a second dose of 2.5 mls (60 mg) 4 to 6 hours after the first dose.3. Give a third dose of 2.5 mls (60 mg) 4 to 6 hours after the second dose.4. Give a fourth dose 4-6 hours after the third dose if your baby still has a fever.
<p>Soreness, swelling* and redness in the area where the injection was given</p>	<ul style="list-style-type: none">• Give paracetamol to relieve aches and pains.• Make sure clothes are not too tight or rubbing against the area where the injection was given.
<p>Mild diarrhoea after the rotavirus vaccine</p>	<ul style="list-style-type: none">• Give extra milk to drink• Wash your hands carefully after changing and disposing of your baby's nappy.

Common reaction	What to do
Fever (over 39°C)	<ul style="list-style-type: none"> • Do not overdress your baby • Make sure their room isn't too hot • Give extra fluids to drink • Give paracetamol to lower the fever
Headache or irritability	<ul style="list-style-type: none"> • Give paracetamol to relieve aches and pains.

* In school children this swelling can be from the shoulder to the elbow.

This usually occurs within 2 days of the vaccination and gets better over 4 – 5 days. Antibiotics are not needed to treat this local reaction.

-
- The dose of paracetamol recommended for your child is written on the bottle according to the child's age.
 - Please ask your pharmacist for sugar-free paracetamol suitable for your child's age.
 - Using paracetamol over a long period without advice from a doctor may be harmful.

For further information

Please read the booklet 'Your child's immunisation - a guide for parents.'

Visit the HSE's immunisation website at www.immunisation.ie

If you are worried about your child please contact your GP, General Practice Nurse, or public health nurse for advice.



Published by: HSE National Immunisation Office

Publication date: May 2023

Reviewed on July 2024

Order code: HNI01679